

Medicine M.C.Q

October 2009

Underline the only one correct answer :

- 1) Pattern of gait in parkinsonism is :  
a- spastic                      b- shuffling                      c- drunken
- 2) One of the following is sign of pyramidal tract lesion except :  
a- hyper-reflexia              b- flexor planter response      c- hypertonia
- 3) Crossed hemiplegia results from a lesion in :  
a- frontal lobe                  b- internal capsule                  c- mid-brain
- 4) Patient with migraine my need :  
a- CT brain                      b- lumbar puncture                  c- none
- 5) Lenticulostriate artery occlusion may cause :  
a- capsular hemiplegia          b- LL monoplegia                  c- paraplegia
- 6) Normal glucose level in CSF occurs in meningitis duo to :  
a- viral                              b- fungal                              c- tuberculosis
- 7) The following are affected in subacute combined degeneration except :  
a- extra pyramidal tract      b- posterior column      c- peripheral nerves
- 8) The following are features of pseudo bulber palsy except :  
a- exaggerated jaw reflex      b- tongue atrophy                  c- nasal tone
- 9) Radicular sensory loss results from a lesion in :  
a- conus medullaries      b- posterior column                  c- cauda equina
- 10) Acute myocardial infarction may be complicated by :  
a- aortic regurge                  b- mitral regurge                  c- pulmonary incompetence
- 11) Atrio-ventricular conduction disturbance does Not occur in :  
a- infectious mononucleosis      b- athletes                              c-  $\alpha$  blockers

البلاء يقرب بينك وبين الله ويعلمك الدعاء ويذهب عبك الفخر والغرور



- 12) Dropped beat occur in :  
a- 1<sup>st</sup> degree HB      b- 2<sup>nd</sup> degree HB      c- 3<sup>rd</sup> degree HB
- 13) Diagnosis of acute myocardial infarction is based on :  
a- chest x-ray      b- CK MB iso-enzyme      c- amylase enzyme
- 14) ECG finding in hyperkalaemia is :  
a- low-amplitude T-wave      b- high-amplitude T-wave      c- U-wave
- 15) Orthopnoea is caused by :  
a- Rt sided heart failure      b- Lt sided heart failure      c- neither
- 16) Agonizing acute chest pain occurs in :  
a- pulmonary embolism      b- anemia      c- heart failure
- 17) Biphasic P-wave in lead II or V1 occurs duo to abnormality of :  
a- Lt ventricle      b- Lt atrium      c- Rt ventricle
- 18) Cardiac cachexia may occur duo to except :  
a- anorexia      b- low metabolic rate      c- protein losing enteropathy
- 19) In paroxysmal supraventricular tachycardia, the rhythm is :  
a- regular      b- regular irregularity      c- irregular regularity
- 20) Causes of prominent R wave in lead V1 :  
a- LV hypertrophy      b- Lt BBB      c- Posterior wall MI
- 21) The commonest cause of constrictive pericarditis is :  
a- rheumatic fever      b- TB      c- viral infection
- 22) P-R interval in wenckback's phenomenon is :  
a- constantly prolonged      b- shortend      c- progressively prolonged
- 23) Main antibody in bronchial asthma is :  
a- IgG      b- IgM      c- IgE
- 24) Sarcoidosis occurs duo to defective :  
a- humeral imunity      b- cell-mediated      c- both

نُيقِنُ أَنَّ الدُّنْيَا دَارُ مَحَنٍ وَبِلَاءٍ وَمُنْغَصَّاتٍ فَاقْبَلْهَا عَلَيَّ حَالَهَا وَاسْتَعْنِ بِاللَّهِ



- 25) Tracheal deviation occurs in except :  
 a- sarcoidosis      b- acute pneumonia      c- pleural effusion
- 26) Alveolo-capillary block may occur in :  
 a- COPD      b- interstitial lung disease      c- both
- 27) In bronchial asthma :  
 a- reduced FEV1/VC      b- increased Pco2      c- increased VC
- 28) Anaphylaxis is :  
 a- type-1 hypersensitivity      b- type-II      c- type-III
- 29) Recognition molecules in immune system are :  
 a- B-cell antigen receptor      b- major histocompatibility      c- both
- 30) One of the following is Not considered a sign of haemolytic anemia :  
 a- pallor      b- splenomegaly      c- itching marking
- 31) Signs of vitamin B12 deficiency anemia include :  
 a- peripheral neuropathy      b- tremors      c- diarrhea
- 32) Skin lesions may occur in Henoch schonlein purpura :  
 a- vesiculobullous      b- pustular      c- both
- 33) In chronic iron deficiency anemia :  
 a- high serum ferritin      b- high TIBC      c- reticulocytosis
- 34) Bilharziasis is treated by :  
 a- praziquantel      b- metronidazole      c- colchicine
- 35) Treatment of acute gouty arthritis is :  
 a- colchicine      b- alluprinol      c- paracetamol
- 36) Pathergy test is helpful in :  
 a- ankylosing spondylitis      b- Behcet's disease      c- scleroderma
- 37) Monoarthritis is a feature of :  
 a- SLE      b- rheumatoid arthritis      c- septic arthritis
- 38) Assessment of disease activity in rheumatoid arthritis depends on except :

اعلم أن من اغتابك فقد أهدي لك حسناؤه وخط من سبناك وجعلك مشهورا وهذه

نعمة



- a- pain scale    b- duration of morning stiffness    c- joint deformity
- 39) The following are non organ specific autoimmune disease except :  
 a- pernicious anemia    b- SLE    c- rheumatoid arthritis
- 40) The following are associated with rheumatoid arthritis except :  
 a- osteoporosis    b- pulmonary fibrosis    c- dementia
- 41) Hypothyroidism is characterized by :  
 a- weight loss    b- bradycardia    c- tremors
- 42) Cancer colon is a common complication in :  
 a- crohn's disease    b- ulcerative colitis    c- chronic amebiasis
- 43) Ulcerative colitis :  
 a- common in smokers    b- common in jejunum    c- associated with HLA27
- 44) The commonest cause of marked rise of liver enzyme is :  
 a- hepatitis A    b- hepatitis B    c- hepatitis C
- 45) Irritable bowel syndrome :  
 a- unrelated to stress    b- pass mucus    c- needs gluten free diet
- 46) Common complication of crhon's disease is :  
 a- peri-anal fistula    b- hepatomegaly    c- cancer colon
- 47) Gastric cancer is commonly :  
 a- squamous cell    b- adenocarcinoma with helicobacter  
 c- cause haematemesis
- 48) Helicobacter pylori :  
 a- is gram +ve bacteria    b- occurs in overcrowding  
 c- cause food poisoning
- 49) Minimal change nephropathy may develop from :  
 a- NSAIDS    b- lithium    c- acyclovir
- 50) Polyuria with high specific gravity occurs in :  
 a- diabetes mellitus    b- diabetes insipidus    c- both

احذر كلمة (سوف) وتأخير الأعمال فهذا عنوان الفشل